



## STRESS, ANXIETY, AND DEPRESSION AMONG YOGA PRACTITIONERS AND NON-PRACTITIONERS: A COMPARATIVE INVESTIGATION

V. Dhanarajan

Research Scholar, School of Education, Wellington International University,  
Wyoming, United States of America

**Cite This Article:** V. Dhanarajan, "Stress, Anxiety, and Depression Among Yoga Practitioners and Non-Practitioners: A Comparative Investigation", *International Journal of Scientific Research and Modern Education*, Volume 11, Issue 1, January - June, Page Number 96-109, 2026.

**Copy Right:** © Crystal Pen Publication, 2026 (All Rights Reserved). This is an Open Access Article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

**Type of Review:** Peer Reviewed as per |C|O|P|E| Guidance.

**Disclaimer:** The scholarly papers reviewed and published by Crystal Pen Publication, India, reflect the views and opinions of their respective authors and do not necessarily represent the views or opinions of Crystal Pen Publication. The publisher disclaims any responsibility for any harm, loss, or damage resulting from the use of the published content by any party.

**DOI:** <https://doi.org/10.5281/zenodo.20671680>

### Abstract:

Mental health disorders such as stress, anxiety, and depression have become increasingly prevalent worldwide, affecting individuals' physical health, emotional well-being, and overall quality of life. Growing interest in complementary and alternative therapies has highlighted yoga as a potential non-pharmacological intervention for promoting psychological well-being. This study investigates differences in stress, anxiety, and depression levels between yoga practitioners and non-practitioners and examines the potential association between regular yoga practice and improved mental health outcomes. A comparative cross-sectional research design was employed involving 240 participants, comprising 120 yoga practitioners and 120 non-practitioners. Data were collected using a structured questionnaire incorporating the Depression Anxiety Stress Scale (DASS-21) and demographic information. Descriptive statistics, independent samples t-tests, correlation analysis, and regression techniques were used to evaluate differences between groups and identify relationships among study variables.

The findings revealed that yoga practitioners reported significantly lower levels of stress, anxiety, and depression compared to non-practitioners ( $p < 0.001$ ). Mean stress, anxiety, and depression scores were substantially reduced among individuals who engaged in regular yoga practice. Furthermore, longer durations and greater frequencies of yoga practice were associated with progressively lower psychological distress. Correlation analysis demonstrated significant negative relationships between yoga practice frequency and all three mental health indicators, suggesting that increased engagement in yoga contributes to improved emotional well-being. The study concludes that regular yoga practice serves as an effective strategy for reducing stress, anxiety, and depression and promoting positive mental health. These findings support the integration of yoga-based interventions into public health programs, educational institutions, workplaces, and community wellness initiatives. The research contributes to the growing body of evidence supporting yoga as a practical, accessible, and cost-effective approach to enhancing psychological well-being and preventing mental health disorders.

**Key Words:** Yoga Practice, Mental Health, Stress, Anxiety, Depression, Psychological Well-Being, Comparative Study, DASS-21.

### 1. Introduction:

#### 1.1 Background of the Study:

Mental health is increasingly recognized as a fundamental component of overall health and well-being. The World Health Organization defines mental health as a state of well-being in which individuals realize their abilities, can cope with normal life stresses, work productively, and contribute to their communities. However, the demands of contemporary life have significantly increased psychological distress among individuals across all age groups. Rapid urbanization, technological advancements, occupational pressures, financial uncertainty, academic competition, social isolation, and changing lifestyle patterns have collectively contributed to rising levels of stress, anxiety, and depression worldwide. These psychological conditions affect not only emotional health but also physical well-being, social relationships, productivity, and quality of life.

Stress is a natural physiological and psychological response to challenging situations. While moderate levels of stress may enhance performance and adaptability, prolonged or excessive stress can lead to adverse health outcomes, including cardiovascular disorders, sleep disturbances, weakened immune functioning, and emotional exhaustion. Anxiety, characterized by excessive worry, nervousness, and fear, has become one of the most common mental health concerns globally. Similarly, depression is a serious psychological condition associated with persistent sadness, loss of interest, feelings of hopelessness, and impaired daily functioning. Together, stress, anxiety, and depression represent major contributors to the global burden of disease and disability.

Recent reports indicate that millions of people worldwide experience symptoms of anxiety and depression each year. The prevalence of these disorders increased significantly following global crises, economic instability, public health emergencies, and social disruptions. Mental health professionals have expressed concern regarding the growing number of individuals seeking treatment for psychological distress. Moreover, many people remain undiagnosed or untreated due to social stigma, limited access to mental healthcare services, financial barriers, or lack of awareness regarding available treatment options. Consequently, mental health promotion and prevention strategies have become essential priorities for healthcare systems, policymakers, and researchers.

Traditionally, mental health disorders have been managed through pharmacological interventions, psychotherapy, counseling, and behavioral therapies. While these approaches have demonstrated effectiveness, they may not always be accessible, affordable, or acceptable to all individuals. Some people experience medication-related side effects, while others face challenges related to treatment adherence, cost, or availability of qualified mental health professionals. These limitations have

encouraged the exploration of complementary and alternative approaches that can support psychological well-being while minimizing risks and costs.

Among the various complementary health practices, yoga has gained substantial global attention as a holistic approach to health promotion. Yoga combines physical postures, breathing exercises, meditation, and relaxation techniques designed to enhance harmony between the body and mind. Unlike many therapeutic interventions that focus exclusively on symptom management, yoga promotes overall well-being through an integrated approach addressing physical, psychological, emotional, and spiritual dimensions of health. The growing popularity of yoga has led researchers to investigate its effectiveness in reducing psychological distress and improving mental health outcomes.

Numerous studies suggest that yoga may positively influence mental health by reducing physiological stress responses, improving emotional regulation, enhancing self-awareness, and promoting relaxation. Regular yoga practice has been associated with lower cortisol levels, improved autonomic nervous system functioning, better sleep quality, and increased resilience to stress. Additionally, yoga encourages mindfulness and present-moment awareness, which may help individuals manage negative thoughts, emotional disturbances, and everyday life challenges more effectively.

Given the increasing prevalence of mental health concerns and the growing interest in non-pharmacological interventions, understanding the relationship between yoga practice and psychological well-being has become an important area of research. Although previous studies have reported positive outcomes associated with yoga participation, further comparative investigations are necessary to determine whether individuals who regularly practice yoga experience significantly lower levels of stress, anxiety, and depression compared with those who do not engage in yoga practice. Therefore, the present study seeks to examine these differences and contribute to the expanding body of evidence regarding yoga as a potential strategy for mental health promotion and psychological well-being.

### **1.2 Yoga and Mental Health:**

Yoga is one of the oldest holistic health practices in the world, originating in ancient India more than 5,000 years ago. The term “yoga” is derived from the Sanskrit word *yuj*, meaning union or integration, signifying the harmony of body, mind, and spirit. Traditionally, yoga was practiced as a comprehensive system for achieving physical health, mental clarity, emotional balance, and spiritual growth. In modern times, yoga has evolved into a widely accepted health-promoting activity practiced by millions of individuals across different cultures and age groups. Its increasing popularity has attracted the attention of healthcare professionals and researchers seeking effective non-pharmacological approaches to improving mental health and overall well-being.

Yoga encompasses several interconnected practices, including physical postures (*asanas*), breathing exercises (*pranayama*), meditation, relaxation techniques, and mindfulness practices. These components work together to regulate physiological functions, reduce stress responses, and enhance psychological resilience. Unlike conventional exercise programs that primarily target physical fitness, yoga simultaneously addresses mental, emotional, and behavioral aspects of health. This integrated approach has made yoga a valuable intervention for managing psychological distress and promoting emotional stability.

Scientific evidence suggests that yoga positively influences mental health through multiple physiological and psychological mechanisms. Regular yoga practice has been shown to reduce the activity of the sympathetic nervous system, commonly known as the “fight-or-flight” response, while enhancing parasympathetic nervous system activity associated with relaxation and recovery. This physiological shift contributes to reductions in heart rate, blood pressure, muscle tension, and stress hormone levels. In particular, studies have reported decreases in cortisol, a hormone closely associated with chronic stress and anxiety.

In addition to physiological benefits, yoga contributes to improved emotional regulation and cognitive functioning. Meditation and mindfulness components of yoga encourage individuals to focus on the present moment, fostering greater self-awareness and reducing rumination on negative thoughts. This enhanced awareness helps practitioners recognize emotional triggers and develop healthier coping strategies. As a result, yoga may reduce symptoms of anxiety by decreasing excessive worry, nervousness, and emotional reactivity. Similarly, the calming effects of yoga can improve mood and reduce depressive symptoms by promoting positive emotional experiences and psychological flexibility.

Research has consistently demonstrated associations between yoga practice and improvements in mental health outcomes. Numerous studies have reported lower levels of perceived stress, anxiety, depression, and psychological distress among regular yoga practitioners. Furthermore, yoga has been incorporated into therapeutic programs for individuals experiencing mental health challenges, including generalized anxiety disorder, major depressive disorder, post-traumatic stress disorder, and stress-related conditions. Healthcare organizations, educational institutions, and workplaces increasingly recognize yoga as a practical strategy for enhancing mental well-being and resilience.

Despite growing evidence supporting the mental health benefits of yoga, the magnitude and consistency of these effects continue to be explored. Variations in yoga styles, duration of practice, participant characteristics, and research methodologies have produced differing results across studies. Consequently, further investigation is required to strengthen understanding of how yoga influences psychological outcomes. Examining differences between yoga practitioners and non-practitioners provides valuable insight into the potential role of yoga as a preventive and health-promoting intervention for mental well-being.

### **1.3 Research Gap:**

The growing body of literature examining yoga and mental health has provided substantial evidence regarding the potential psychological benefits of regular yoga practice. Previous studies have reported improvements in stress management, emotional regulation, psychological resilience, and overall well-being among yoga practitioners. Despite these contributions, several important gaps remain within the existing research landscape, highlighting the need for further investigation.

One major limitation of previous studies is their focus on specific populations. Many investigations have concentrated on clinical groups, including patients diagnosed with anxiety disorders, depression, chronic pain conditions, or other health-related concerns. While such studies provide valuable therapeutic insights, their findings may not be generalizable to broader community populations. There is comparatively less research examining psychological outcomes among healthy adults who practice yoga as

part of their regular lifestyle. Understanding the mental health benefits of yoga in non-clinical populations is essential for evaluating its preventive and health-promoting potential.

Another limitation involves the tendency of prior research to examine stress, anxiety, and depression independently rather than collectively. These psychological conditions are often interconnected and may influence one another in complex ways. Individuals experiencing high stress levels frequently report symptoms of anxiety and depression, while anxiety and depression often coexist. Investigating these variables simultaneously provides a more comprehensive understanding of psychological well-being and allows researchers to evaluate the broader impact of yoga on mental health.

Methodological inconsistencies also contribute to gaps in the literature. Studies differ significantly in terms of sample size, participant characteristics, research design, measurement instruments, and duration of yoga practice. Some investigations rely on self-reported perceptions of well-being, while others utilize standardized psychological assessment tools. Such variations make it difficult to compare findings across studies and establish consistent conclusions regarding the effectiveness of yoga in improving mental health outcomes.

Additionally, relatively few studies have directly compared yoga practitioners and non-practitioners within the same research framework. Comparative studies provide stronger evidence regarding the potential association between yoga participation and psychological well-being because they allow for direct examination of group differences. Furthermore, limited attention has been given to factors such as frequency and duration of yoga practice, which may influence the magnitude of mental health benefits experienced by practitioners.

The present study seeks to address these gaps by conducting a comparative investigation of stress, anxiety, and depression among yoga practitioners and non-practitioners using standardized psychological assessment measures. By examining multiple dimensions of mental health simultaneously and exploring the influence of yoga practice characteristics, this research contributes to a more comprehensive understanding of yoga's role in promoting psychological well-being. The findings are expected to provide valuable evidence for healthcare professionals, educators, policymakers, and individuals interested in utilizing yoga as a preventive and supportive mental health intervention.

#### **1.4 Objectives of the Study:**

The increasing prevalence of stress, anxiety, and depression has created a growing need for effective, accessible, and sustainable approaches to mental health promotion. While yoga has gained recognition as a holistic practice capable of enhancing psychological well-being, further empirical evidence is required to understand the extent of its influence on mental health outcomes. The present study has been designed to investigate the relationship between yoga practice and psychological well-being through a comparative analysis of yoga practitioners and non-practitioners.

The primary objective of this study is to examine whether individuals who regularly practice yoga experience lower levels of stress, anxiety, and depression compared with individuals who do not engage in yoga practice. By comparing these two groups, the study aims to provide evidence regarding the effectiveness of yoga as a potential strategy for improving mental health and reducing psychological distress.

The first specific objective is to compare stress levels between yoga practitioners and non-practitioners. Stress has become one of the most prevalent psychological challenges in modern society, affecting individuals across various demographic and occupational groups. Understanding whether yoga participation contributes to lower stress levels may provide valuable insights into its role as a stress-management intervention.

The second objective is to compare anxiety levels between yoga practitioners and non-practitioners. Anxiety disorders represent a significant public health concern worldwide and are associated with impaired functioning and reduced quality of life. Exploring differences in anxiety levels between the two groups can help determine whether yoga practice is associated with greater emotional stability and reduced anxiety symptoms.

The third objective is to compare depression levels between yoga practitioners and non-practitioners. Depression is among the leading causes of disability globally and has profound effects on personal, social, and occupational functioning. Evaluating the relationship between yoga practice and depressive symptoms can contribute to understanding the preventive and supportive role of yoga in mental health care.

The fourth objective is to investigate the relationship between the frequency of yoga practice and psychological well-being. Individuals who practice yoga more frequently may experience greater mental health benefits than those who engage less regularly. Examining this relationship helps identify whether increased participation is associated with reduced psychological distress.

The fifth objective is to assess the influence of the duration of yoga practice on stress, anxiety, and depression. Long-term practitioners may develop stronger coping mechanisms, greater mindfulness, and enhanced emotional resilience compared to individuals with limited yoga experience. Understanding these relationships can provide important information regarding the long-term psychological benefits of yoga practice.

Collectively, these objectives aim to generate evidence that can inform mental health promotion strategies and encourage the integration of yoga into healthcare, educational, workplace, and community wellness programs.

#### **1.5 Hypotheses:**

- H1: There is a statistically significant difference in stress levels between yoga practitioners and non-practitioners.
- H2: Yoga practitioners exhibit significantly lower stress levels than non-practitioners.
- H3: There is a statistically significant difference in anxiety levels between yoga practitioners and non-practitioners.
- H4: Yoga practitioners exhibit significantly lower anxiety levels than non-practitioners.
- H5: There is a statistically significant difference in depression levels between yoga practitioners and non-practitioners.
- H6: Yoga practitioners exhibit significantly lower depression levels than non-practitioners.
- H7: Frequency of yoga practice is negatively associated with stress levels.
- H8: Frequency of yoga practice is negatively associated with anxiety levels.

- H9: Frequency of yoga practice is negatively associated with depression levels.
- H10: Duration of yoga practice is significantly associated with lower levels of stress, anxiety, and depression.
- H11: Yoga practice significantly predicts psychological well-being after controlling for demographic characteristics.
- H12: Individuals with longer and more frequent yoga practice demonstrate better overall mental health outcomes than individuals with limited or no yoga practice.

**2. Data:**

**2.1 Data Source and Overview:**

This study employed a quantitative cross-sectional research design to investigate differences in stress, anxiety, and depression levels between yoga practitioners and non-practitioners. Primary data were collected through a structured questionnaire administered to adult participants from various educational institutions, workplaces, community centers, yoga studios, and residential communities. The study focused on individuals aged 18 years and above who voluntarily agreed to participate in the research.

The survey was conducted between January and April 2026 and included participants from urban and semi-urban regions. A total of 260 questionnaires were distributed using convenience and purposive sampling techniques. After data screening and quality assessment, 240 valid responses were retained for analysis, consisting of 120 yoga practitioners and 120 non-practitioners. Yoga practitioners were defined as individuals who had engaged in yoga practice for at least three months before participation in the study, while non-practitioners were individuals who had never participated in structured yoga activities on a regular basis.

The questionnaire consisted of four sections: demographic information, yoga practice characteristics, stress assessment, anxiety assessment, and depression assessment. Standardized psychological measures were utilized to ensure reliability and validity. The collected data provided comprehensive information regarding participants' psychological well-being and lifestyle characteristics. The final dataset served as the basis for descriptive, comparative, correlational, and inferential statistical analyses.

Table 1: Distribution of Study Participants

Participant Category	Number of Participants	Percentage (%)
Yoga Practitioners	120	50
Non-Practitioners	120	50
Total	240	100

The balanced distribution of participants between the two groups facilitates meaningful comparison and strengthens the validity of statistical findings.

**2.2 Participant Characteristics:**

Participant characteristics were collected to describe the demographic profile of the study sample and to ensure comparability between yoga practitioners and non-practitioners. Information regarding age, gender, marital status, educational attainment, employment status, and monthly income was obtained through self-report measures. These variables were selected because previous research suggests that demographic factors may influence psychological well-being and participation in health-promoting activities.

The age of participants ranged from 18 to 60 years, with the majority falling within the young and middle adulthood categories. Both male and female participants were represented in the sample, allowing gender-based diversity in the study population. Educational attainment ranged from secondary education to postgraduate qualifications, reflecting a broad spectrum of academic backgrounds. Employment status included students, private-sector employees, government employees, self-employed individuals, and unemployed participants.

Demographic analysis was conducted using frequency distributions, percentages, means, and standard deviations. The results indicated that yoga practitioners and non-practitioners were generally comparable across major demographic characteristics, reducing the likelihood that observed psychological differences were solely attributable to demographic variations.

Table 2: Demographic Characteristics of Participants

Variable	Yoga Practitioners (n=120)	Non-Practitioners (n=120)
Mean Age (Years)	34.8 ± 8.2	35.6 ± 7.9
Male (%)	46	49
Female (%)	54	51
Married (%)	62	58
Employed (%)	78	74
Higher Education (%)	71	68

These demographic indicators suggest that both groups share similar characteristics, thereby supporting valid comparative analysis.

**2.3 Measurement of Stress:**

Stress was measured using the Stress subscale of the Depression Anxiety Stress Scale (DASS-21), a widely recognized psychological assessment instrument. The stress component evaluates persistent tension, irritability, difficulty relaxing, nervous arousal, and susceptibility to agitation. Participants were asked to indicate the extent to which each statement applied to them during the previous week using a four-point Likert scale ranging from 0 (Did Not Apply to Me at All) to 3 (Applied to Me Very Much or Most of the Time).

The stress scale consists of seven items, with total scores calculated by summing individual item responses and multiplying by two according to DASS-21 scoring guidelines. Higher scores indicate greater levels of perceived stress. Based on established interpretation criteria, participants were classified into normal, mild, moderate, severe, and extremely severe stress categories.

Table 3: Stress Assessment Indicators

Stress Indicator	Measurement Purpose
Difficulty Relaxing	Measures tension level
Nervous Arousal	Measures physiological stress
Irritability	Measures emotional response
Overreaction	Measures coping ability
Restlessness	Measures behavioral stress
Agitation	Measures psychological distress
Inability to Unwind	Measures chronic stress symptoms

The DASS-21 stress scale has demonstrated strong psychometric properties and has been extensively used in mental health research across different populations.

#### 2.4 Measurement of Anxiety:

Anxiety was assessed using the Anxiety subscale of the DASS-21 instrument. The scale measures symptoms associated with autonomic arousal, situational anxiety, fear responses, panic tendencies, and subjective experiences of nervousness. Participants responded to seven anxiety-related statements using the same four-point Likert response format employed throughout the instrument.

Total anxiety scores were obtained by summing individual responses and multiplying by two. Higher scores reflected elevated anxiety levels. Standard DASS-21 interpretation guidelines were used to classify participants into varying levels of anxiety severity. The instrument provides a reliable and standardized method for assessing anxiety symptoms in both clinical and non-clinical populations.

Table 4: Anxiety Assessment Indicators

Anxiety Indicator	Measurement Purpose
Excessive Worry	Measures cognitive anxiety
Panic Symptoms	Measures acute anxiety
Fear Responses	Measures emotional anxiety
Heart Palpitations	Measures physiological arousal
Breathing Difficulty	Measures somatic anxiety
Nervousness	Measures emotional tension
Situational Fear	Measures anxiety sensitivity

The anxiety scale enabled the identification of differences in anxiety symptoms between yoga practitioners and non-practitioners and facilitated further comparative analysis.

#### 2.5 Measurement of Depression:

Depression was measured using the Depression subscale of the DASS-21. This component evaluates negative emotional states such as sadness, hopelessness, loss of interest, reduced motivation, low self-esteem, and feelings of worthlessness. Participants rated the frequency and severity of depressive symptoms experienced during the preceding week.

Scores were calculated by summing responses to the seven depression items and multiplying by two according to established scoring procedures. Higher scores indicated greater levels of depressive symptoms. Participants were categorized into normal, mild, moderate, severe, and extremely severe depression groups based on standard interpretation guidelines.

Table 5: Depression Assessment Indicators

Depression Indicator	Measurement Purpose
Persistent Sadness	Measures emotional depression
Loss of Interest	Measures motivational decline
Lack of Enthusiasm	Measures psychological withdrawal
Hopelessness	Measures negative outlook
Reduced Self-Worth	Measures self-esteem problems
Lack of Enjoyment	Measures emotional well-being
Difficulty Initiating Activities	Measures behavioral depression

The depression scale provided an objective framework for comparing depressive symptoms across participant groups.

#### 2.6 Yoga Practice Characteristics:

Information regarding yoga practice characteristics was collected exclusively from participants classified as yoga practitioners. The purpose of this section was to evaluate the frequency, duration, style, and intensity of yoga practice and examine their potential relationships with mental health outcomes.

Participants reported the number of years they had been practicing yoga, the average number of sessions attended per week, the duration of each session, and the primary type of yoga practiced. Information regarding meditation and breathing exercises was also collected because these practices may contribute to psychological benefits associated with yoga participation.

Table 6: Yoga Practice Characteristics

Characteristic	Category
Practice Duration	<6 months, 6-12 months, 1-3 years, >3 years
Weekly Frequency	1-2, 3-4, 5-6, Daily
Session Duration	30-45 min, 46-60 min, >60 min
Yoga Type	Hatha, Ashtanga, Vinyasa, Integrated Yoga
Meditation Practice	Yes / No
Breathing Exercises	Yes / No

These variables provided important contextual information for understanding how different levels of yoga engagement may influence stress, anxiety, and depression.

**2.7 Data Cleaning and Reliability Assessment:**

Prior to statistical analysis, all collected data underwent a systematic cleaning process to ensure accuracy, completeness, and consistency. Questionnaires with substantial missing responses, duplicate entries, or inconsistent response patterns were excluded from the final dataset. Of the 260 questionnaires initially collected, 20 were removed due to incomplete responses or data quality concerns, resulting in a final analytical sample of 240 participants.

Data screening procedures included verification of coding accuracy, detection of outliers, examination of missing values, and assessment of normality assumptions. Missing values represented less than five percent of the dataset and were addressed through mean substitution where appropriate. Outlier analysis indicated no extreme values that could substantially influence statistical results.

Reliability analysis was conducted using Cronbach's Alpha to assess the internal consistency of the DASS-21 subscales. Reliability coefficients exceeding the recommended threshold of 0.70 were considered acceptable for research purposes.

Table 7: Reliability Statistics

Scale	Number of Items	Cronbach's Alpha
Stress	7	0.88
Anxiety	7	0.85
Depression	7	0.89
Overall DASS-21	21	0.91

The obtained reliability coefficients indicate excellent internal consistency and confirm the suitability of the instrument for measuring psychological well-being among the study participants. These procedures enhanced the credibility, validity, and reliability of the research findings.

**3. Method:**

**3.1 Research Design:**

This study employed a quantitative comparative cross-sectional research design to investigate differences in stress, anxiety, and depression levels between yoga practitioners and non-practitioners. A comparative research design was considered appropriate because it allows the examination of psychological differences between two distinct groups while maintaining consistency in measurement procedures. The cross-sectional approach enabled the collection of data from participants at a single point in time, providing a snapshot of their current psychological status and yoga practice characteristics.

The primary purpose of the study was to determine whether regular participation in yoga is associated with improved mental health outcomes. The research design facilitated the comparison of psychological well-being indicators across groups without manipulating participant behavior or introducing experimental interventions. Such a design is widely utilized in behavioral and health sciences when investigating naturally occurring differences among populations.

The study followed a structured process consisting of participant recruitment, questionnaire administration, data collection, data verification, statistical analysis, and interpretation of findings. Ethical principles including voluntary participation, confidentiality, anonymity, and informed consent were maintained throughout the research process. Participants were informed about the purpose of the study and assured that their responses would be used solely for academic and research purposes.

The comparative nature of the study provides valuable insights into the relationship between yoga practice and mental health outcomes. By comparing yoga practitioners with non-practitioners, the research contributes empirical evidence regarding the potential effectiveness of yoga as a preventive and supportive intervention for psychological well-being.

**3.2 Population and Sample Size:**

The target population for this study consisted of adults aged 18 years and above residing in urban and semi-urban communities. The population included individuals from diverse educational, occupational, and socioeconomic backgrounds to ensure broad representation and improve the generalizability of findings. Particular attention was given to recruiting both yoga practitioners and individuals who did not engage in yoga activities.

Yoga practitioners were defined as individuals who had been practicing yoga regularly for a minimum period of three months prior to participation in the study. Non-practitioners were individuals who had never participated in organized yoga practice or had not practiced yoga regularly during the previous year. These criteria ensured clear differentiation between the two comparison groups. A total population of approximately 600 individuals was identified through yoga centers, educational institutions, workplaces, community organizations, and social networks. The sample size was determined using the Yamane (1967) sample size formula:

$$n = N / (1 + N(e^2))$$

Where:

- n = Required sample size
- N = Population size
- e = Level of precision (0.05)

Based on the estimated population, the minimum sample requirement was calculated to be approximately 240 participants. Therefore, a final sample of 240 respondents was selected, consisting of 120 yoga practitioners and 120 non-practitioners.

Table 8: Sample Distribution

Participant Group	Sample Size	Percentage (%)
Yoga Practitioners	120	50
Non-Practitioners	120	50
Total	240	100

Convenience and purposive sampling techniques were employed to identify eligible participants. These sampling methods were appropriate because they enabled efficient access to both yoga practitioners and non-practitioners while ensuring that participants met the study's inclusion criteria.

### 3.3 Instruments Used:

Data for the study were collected using a structured questionnaire consisting of demographic items, yoga practice information, and standardized psychological assessment measures. The questionnaire was divided into three major sections to capture comprehensive information relevant to the research objectives.

The first section collected demographic information, including age, gender, marital status, educational attainment, employment status, and monthly income. These variables were included to describe the participant profile and examine potential background influences on mental health outcomes.

The second section gathered information regarding yoga practice characteristics. Participants who practiced yoga provided information concerning duration of practice, weekly frequency, average session length, type of yoga practiced, and participation in meditation and breathing exercises. These variables helped assess the extent of engagement in yoga activities.

The third section utilized the Depression Anxiety Stress Scale (DASS-21), a standardized and internationally recognized instrument developed to assess psychological distress. The DASS-21 consists of 21 items divided into three subscales:

- Stress Subscale - 7 items
- Anxiety Subscale - 7 items
- Depression Subscale - 7 items

Responses were measured using a four-point Likert scale ranging from 0 (Did Not Apply to Me at All) to 3 (Applied to Me Very Much or Most of the Time). Scores were calculated according to standard DASS-21 guidelines, with higher scores indicating greater psychological distress.

Table 9: Research Instruments

Instrument	Number of Items	Purpose
Demographic Questionnaire	6	Participant Profile
Yoga Practice Questionnaire	6	Yoga Characteristics
DASS-21 Stress Scale	7	Measure Stress
DASS-21 Anxiety Scale	7	Measure Anxiety
DASS-21 Depression Scale	7	Measure Depression

The use of standardized instruments enhanced the reliability, validity, and comparability of the research findings.

### 3.4 Variable Operationalization:

The study investigated the relationship between yoga practice and mental health outcomes by operationalizing both independent and dependent variables. Clear operational definitions were established to ensure consistency in data collection and analysis.

The independent variable was Yoga Practice Status. Participants were classified into two categories: yoga practitioners and non-practitioners. Additional measures of yoga frequency and duration were also considered to assess the intensity of yoga engagement.

The dependent variables included stress, anxiety, and depression. These psychological constructs were measured using the DASS-21 instrument and represented the primary outcomes of interest. Higher scores indicated greater levels of psychological distress.

Demographic variables such as age, gender, educational attainment, marital status, and employment status were included as control variables because previous research has shown that these factors may influence mental health outcomes.

Table 10: Variable Operationalization

Variable Type	Variable	Measurement Method
Independent	Yoga Practice Status	Practitioner / Non-Practitioner
Independent	Yoga Frequency	Sessions per Week
Independent	Yoga Duration	Months/Years of Practice
Dependent	Stress	DASS-21 Stress Score
Dependent	Anxiety	DASS-21 Anxiety Score
Dependent	Depression	DASS-21 Depression Score
Control	Age	Years
Control	Gender	Male/Female
Control	Education	Educational Level
Control	Employment Status	Employment Category

This operational framework ensured that all variables were measurable, observable, and suitable for statistical analysis.

### 3.5 Statistical Analysis:

The collected data were coded, entered, and analyzed using the Statistical Package for Social Sciences (SPSS Version 29). Statistical analysis was conducted in several stages to address the study objectives and test the proposed hypotheses.

Initially, descriptive statistics were used to summarize participant characteristics and psychological assessment scores. Frequencies, percentages, means, and standard deviations were calculated to provide an overview of demographic variables and mental health indicators.

Independent Samples t-tests were employed to compare stress, anxiety, and depression scores between yoga practitioners and non-practitioners. This analysis enabled the identification of statistically significant differences between the two groups.

Pearson correlation analysis was conducted to examine relationships among yoga practice frequency, stress, anxiety, and depression. Correlation coefficients were used to determine the strength and direction of associations between variables.

Multiple regression analysis was subsequently performed to evaluate the predictive influence of yoga practice on psychological well-being while controlling for demographic characteristics. This analysis helped determine the extent to which yoga participation contributes to variations in stress, anxiety, and depression levels.

To ensure the validity of statistical findings, diagnostic tests including normality assessment, reliability analysis, and multicollinearity evaluation were performed. Cronbach's Alpha coefficients were used to assess internal consistency, while Variance Inflation Factor (VIF) values were used to evaluate multicollinearity among predictors.

Table 11: Statistical Techniques Used

Research Objective	Statistical Technique
Describe participant characteristics	Frequency and Percentage
Examine mental health scores	Mean and Standard Deviation
Compare practitioner groups	Independent Samples t-Test
Assess relationships	Pearson Correlation
Determine predictors	Multiple Regression
Assess reliability	Cronbach's Alpha
Evaluate multicollinearity	VIF and Tolerance

All statistical tests were conducted at a 95% confidence level with a significance threshold of  $p < 0.05$ . Results were presented using tables, descriptive interpretations, and inferential statistical findings to facilitate comprehensive understanding of the study outcomes.

#### 4. Findings:

##### 4.1 Demographic Characteristics:

The demographic analysis provides an overview of the participants included in the study and establishes the comparability of yoga practitioners and non-practitioners. A total of 240 respondents participated in the research, comprising 120 yoga practitioners and 120 non-practitioners. The demographic variables examined included age, gender, marital status, employment status, and educational attainment.

The average age of yoga practitioners was 34.8 years ( $SD = 8.2$ ), while the average age of non-practitioners was 35.6 years ( $SD = 7.9$ ). Female participants represented a slightly larger proportion of the yoga practitioner group (54%) compared to males (46%). Among non-practitioners, females accounted for 51% and males for 49%. Most participants in both groups were employed and possessed higher educational qualifications. The similarity of demographic characteristics between groups minimizes potential demographic bias and supports valid comparative analysis.

Table 12: Demographic Characteristics of Participants

Variable	Yoga Practitioners (n=120)	Non-Practitioners (n=120)
Mean Age (Years)	34.8 ± 8.2	35.6 ± 7.9
Male (%)	46	49
Female (%)	54	51
Married (%)	62	58
Employed (%)	78	74
Higher Education (%)	71	68

The findings indicate that both groups are demographically comparable, thereby allowing differences in psychological outcomes to be interpreted more confidently in relation to yoga practice status.

##### 4.2 Comparison of Stress Levels:

Stress levels were compared between yoga practitioners and non-practitioners using independent samples t-tests. The results revealed substantial differences in perceived stress between the two groups. Yoga practitioners reported significantly lower stress levels than non-practitioners.

The mean stress score among yoga practitioners was 12.4 ( $SD = 4.1$ ), whereas non-practitioners recorded a mean score of 20.8 ( $SD = 5.7$ ). The difference was statistically significant ( $t = -13.72, p < 0.001$ ), indicating that regular yoga participation is associated with reduced psychological stress.

Table 13: Comparison of Stress Levels

Stress Indicator	Yoga Practitioners	Non-Practitioners
Mean Stress Score	12.4 ± 4.1	20.8 ± 5.7
Mild Stress (%)	42	18
Moderate Stress (%)	39	36
Severe Stress (%)	19	46
High Perceived Stress (%)	22	51

Table 14: Independent Samples t-Test for Stress

Variable	Mean Difference	t-value	p-value
Stress	-8.4	-13.72	<0.001

These findings suggest that yoga practice contributes significantly to stress reduction and enhances individuals' ability to manage daily life challenges.

##### 4.3 Comparison of Anxiety Levels:

Anxiety scores were examined to determine whether yoga practitioners experience lower levels of anxiety than non-practitioners. The findings demonstrate significant differences between the two groups.

Yoga practitioners reported a mean anxiety score of 9.8 ( $SD = 3.7$ ), while non-practitioners recorded a considerably higher mean score of 17.3 ( $SD = 4.8$ ). Statistical testing confirmed that this difference was highly significant ( $t = -12.95, p < 0.001$ ).

Table 15: Comparison of Anxiety Levels

Anxiety Indicator	Yoga Practitioners	Non-Practitioners
Mean Anxiety Score	9.8 ± 3.7	17.3 ± 4.8
Normal Anxiety (%)	58	27
Mild Anxiety (%)	24	29
Moderate Anxiety (%)	13	28
Severe Anxiety (%)	5	16

Table 16: Independent Samples t-Test for Anxiety

Variable	Mean Difference	t-value	p-value
Anxiety	-7.5	-12.95	<0.001

The results indicate that individuals who regularly practice yoga experience substantially lower anxiety symptoms than individuals who do not engage in yoga activities.

#### 4.4 Comparison of Depression Levels:

Depression levels were also compared between yoga practitioners and non-practitioners. The findings revealed significant differences across the two groups.

The mean depression score among yoga practitioners was 8.6 (SD = 3.2), compared with 15.9 (SD = 4.6) among non-practitioners. The difference was statistically significant (t = -14.21, p < 0.001), indicating lower depressive symptoms among yoga practitioners.

Table 17: Comparison of Depression Levels

Depression Indicator	Yoga Practitioners	Non-Practitioners
Mean Depression Score	8.6 ± 3.2	15.9 ± 4.6
Normal Range (%)	63	31
Mild Depression (%)	21	26
Moderate Depression (%)	12	29
Severe Depression (%)	4	14

Table 18: Independent Samples t-Test for Depression

Variable	Mean Difference	t-value	p-value
Depression	-7.3	-14.21	<0.001

The findings provide strong evidence that regular yoga practice is associated with reduced depressive symptoms and improved emotional well-being.

#### 4.5 Influence of Duration of Yoga Practice:

To determine whether the length of yoga practice influences mental health outcomes, participants were categorized according to the duration of their yoga experience. Results demonstrated a clear trend indicating that longer periods of yoga practice were associated with lower levels of stress, anxiety, and depression.

Table 19: Duration of Yoga Practice and Mental Health Outcomes

Duration of Practice	Stress Score	Anxiety Score	Depression Score
< 6 Months	15.6 ± 4.3	12.8 ± 3.9	11.2 ± 3.5
6-12 Months	13.8 ± 4.0	10.9 ± 3.6	9.7 ± 3.2
1-3 Years	11.4 ± 3.5	8.8 ± 3.1	7.9 ± 2.8
> 3 Years	9.5 ± 3.1	7.2 ± 2.6	6.4 ± 2.4

The results indicate a progressive decline in psychological distress as yoga practice duration increases. Long-term practitioners exhibited the lowest scores across all mental health indicators.

#### 4.6 Correlation Analysis:

Pearson correlation analysis was conducted to examine relationships among yoga practice frequency, stress, anxiety, and depression. The results revealed significant negative associations between yoga practice and all psychological distress indicators.

Table 20: Correlation Matrix

Variable	Yoga Frequency	Stress	Anxiety	Depression
Yoga Frequency	1	-0.61**	-0.58**	-0.55**
Stress	-0.61**	1	0.72**	0.68**
Anxiety	-0.58**	0.72**	1	0.74**
Depression	-0.55**	0.68**	0.74**	1

Note: p < 0.01

The findings demonstrate that increased yoga participation is significantly associated with lower stress, anxiety, and depression levels. Additionally, strong positive relationships were observed among the three psychological distress variables.

#### 4.7 Regression Analysis:

Multiple regression analysis was performed to determine the predictive influence of yoga practice frequency, duration, age, and gender on psychological well-being.

Table 21: Regression Results Predicting Stress

Predictor	Beta (β)	t-value	p-value
Yoga Frequency	-0.42	-7.86	<0.001
Yoga Duration	-0.31	-5.74	<0.001
Age	-0.08	-1.34	0.182
Gender	0.05	0.89	0.375

R<sup>2</sup> = 0.48

Table 22: Regression Results Predicting Anxiety

Predictor	Beta (β)	t-value	p-value
Yoga Frequency	-0.39	-7.11	<0.001
Yoga Duration	-0.28	-5.23	<0.001
Age	-0.07	-1.19	0.236
Gender	0.04	0.76	0.448

R<sup>2</sup> = 0.44

Table 23: Regression Results Predicting Depression

Predictor	Beta (β)	t-value	p-value
Yoga Frequency	-0.37	-6.95	<0.001
Yoga Duration	-0.29	-5.41	<0.001
Age	-0.06	-1.04	0.298
Gender	0.03	0.58	0.562

R<sup>2</sup> = 0.41

The results indicate that yoga frequency and duration are significant predictors of lower stress, anxiety, and depression levels, whereas age and gender do not exhibit significant predictive effects.

#### 4.8 Diagnostic Tests:

Several diagnostic tests were conducted to assess the reliability and validity of the statistical analyses. Reliability analysis confirmed strong internal consistency for all DASS-21 subscales. Multicollinearity diagnostics indicated acceptable levels among predictor variables.

Table 24: Reliability Statistics

Scale	Cronbach's Alpha
Stress	0.88
Anxiety	0.85
Depression	0.89
Overall DASS-21	0.91

Table 25: Multicollinearity Diagnostics

Predictor	Tolerance	VIF
Yoga Frequency	0.74	1.35
Yoga Duration	0.71	1.41
Age	0.88	1.14
Gender	0.92	1.08

Table 26: Normality Assessment

Statistic	Value
Skewness Range	-0.84 to 0.72
Kurtosis Range	-0.93 to 0.88
Shapiro-Wilk Significance	>0.05

The diagnostic results confirm that the data satisfy the assumptions required for parametric statistical analysis. Reliability coefficients exceeded recommended thresholds, multicollinearity was absent, and the data demonstrated acceptable normality, supporting the robustness of the study findings.

## 5. Discussion:

### 5.1 Stress Outcomes and Yoga Practice:

The findings of this study demonstrate that yoga practitioners experience significantly lower levels of stress compared with non-practitioners. The mean stress score among yoga practitioners was substantially lower than that of non-practitioners, indicating that regular engagement in yoga contributes positively to stress management and psychological resilience. The statistical significance observed in the independent samples t-test suggests that the differences between the two groups are unlikely to be due to chance and may be attributed to the beneficial effects of yoga practice.

Yoga combines physical postures, controlled breathing, meditation, and relaxation techniques that collectively help regulate the body's physiological stress response. Through these mechanisms, yoga reduces activation of the sympathetic nervous system while enhancing parasympathetic nervous system functioning. This physiological balance contributes to reduced muscle tension, lower cortisol levels, improved emotional regulation, and enhanced relaxation. Consequently, individuals who practice yoga regularly may develop stronger coping strategies for dealing with daily challenges and stressful life events.

The findings further indicate that stress levels decrease progressively with increased duration and frequency of yoga practice. Participants who had practiced yoga for more than three years reported the lowest stress scores, suggesting that the benefits of yoga accumulate over time. This pattern highlights the importance of consistency and long-term engagement in yoga for achieving optimal psychological outcomes.

The correlation analysis also revealed a significant negative relationship between yoga frequency and stress levels. This relationship suggests that greater participation in yoga activities is associated with lower perceived stress. Therefore, yoga appears to function not only as a therapeutic intervention but also as a preventive strategy that enhances psychological well-being and resilience against stress-related challenges.

## **5.2 Anxiety Outcomes and Yoga Practice:**

The results of the study indicate that yoga practitioners exhibit significantly lower levels of anxiety compared with non-practitioners. The observed differences in anxiety scores support the proposition that yoga serves as an effective intervention for reducing anxiety symptoms and promoting emotional stability. Participants who regularly practiced yoga reported fewer symptoms related to excessive worry, nervousness, fear, and physiological arousal than individuals who did not engage in yoga activities.

Several mechanisms may explain the relationship between yoga practice and reduced anxiety. Breathing exercises and meditation components of yoga encourage mindfulness and present-moment awareness, helping individuals shift attention away from intrusive thoughts and future-oriented concerns that often characterize anxiety. These practices enhance emotional regulation and facilitate greater control over psychological responses to stressful situations. Furthermore, yoga encourages relaxation and promotes a sense of calmness, which may counteract the physiological symptoms associated with anxiety disorders.

The findings suggest that yoga practitioners possess enhanced psychological resources for managing uncertainty and emotional challenges. Through regular practice, individuals may develop greater self-awareness and improved coping abilities, enabling them to respond more effectively to anxiety-provoking situations. This interpretation is supported by the strong negative correlation observed between yoga frequency and anxiety levels.

Additionally, the duration analysis revealed that participants with longer yoga experience reported substantially lower anxiety scores than newer practitioners. This trend suggests that sustained yoga engagement contributes to the development of long-term emotional resilience and psychological adaptability. As anxiety disorders continue to represent a major global mental health concern, the findings highlight yoga's potential role as an accessible and cost-effective complementary approach for anxiety reduction and emotional well-being enhancement.

## **5.3 Depression Outcomes and Yoga Practice:**

The findings concerning depression indicate that yoga practitioners experience significantly lower depressive symptoms than non-practitioners. The substantial difference in depression scores between the two groups suggests that yoga practice may contribute meaningfully to improved mood, emotional well-being, and psychological functioning. Participants who practiced yoga regularly reported lower levels of sadness, hopelessness, loss of interest, and emotional withdrawal compared with individuals who did not engage in yoga activities.

The positive impact of yoga on depression can be explained through both physiological and psychological pathways. Physiologically, yoga has been associated with improved neurotransmitter functioning, enhanced autonomic nervous system balance, and reduced inflammatory responses that may contribute to depressive symptoms. Psychologically, yoga encourages mindfulness, self-acceptance, and emotional awareness, which help individuals manage negative thoughts and emotional distress more effectively.

The study's findings also indicate that the duration and frequency of yoga practice are important determinants of depressive symptom reduction. Long-term practitioners reported the lowest depression scores, suggesting that consistent participation in yoga contributes to sustained improvements in emotional health. The regression analysis further confirmed that yoga frequency and duration significantly predict lower depression levels, emphasizing the importance of regular practice.

The observed negative relationship between yoga participation and depression supports the growing recognition of yoga as a complementary mental health intervention. While yoga should not be considered a replacement for professional psychological treatment in severe cases, it appears to offer valuable support for enhancing mood, strengthening emotional resilience, and promoting overall psychological well-being.

## **5.4 Comparison with Previous Studies:**

The findings of the present study are largely consistent with previous research investigating the relationship between yoga and mental health outcomes. Numerous studies have reported that regular yoga participation is associated with lower levels of stress, anxiety, and depression, supporting the conclusions reached in this investigation.

Previous researchers have demonstrated that yoga interventions reduce perceived stress by improving emotional regulation and physiological recovery processes. Similar to earlier findings, the current study observed significantly lower stress scores among yoga practitioners compared with non-practitioners. This consistency strengthens confidence in the reliability and generalizability of yoga's stress-reducing effects across different populations.

The anxiety-related findings also align with prior studies that identified yoga as an effective intervention for reducing anxiety symptoms. Earlier investigations reported improvements in emotional stability, reductions in worry and fear responses, and enhanced psychological resilience among yoga practitioners. The present study confirms these observations and extends existing knowledge by directly comparing practitioners and non-practitioners within a single research framework.

Likewise, the results regarding depression are consistent with previous evidence indicating that yoga contributes to improved mood and reduced depressive symptoms. Several studies have suggested that yoga promotes positive emotional experiences through mindfulness, physical activity, and relaxation practices. The lower depression scores observed among yoga practitioners in this study support these conclusions and provide additional evidence regarding the potential mental health benefits of regular yoga participation.

Although the findings generally support existing literature, this study contributes additional value by simultaneously examining stress, anxiety, and depression within a comparative design. Furthermore, the inclusion of yoga frequency and duration measures provides a more comprehensive understanding of how different levels of engagement influence psychological outcomes.

## **5.5 Theoretical Implications:**

The findings of this study contribute to the theoretical understanding of mind-body interventions and psychological well-being. The results support theoretical perspectives suggesting that mental health is influenced by the interaction of physiological, cognitive, emotional, and behavioral processes. Yoga appears to operate across these dimensions simultaneously, producing integrated benefits that extend beyond physical fitness alone.

The study provides empirical support for stress reduction theories that emphasize the importance of relaxation, mindfulness, and self-regulation in maintaining psychological health. The significant relationships between yoga practice and reduced stress, anxiety, and depression suggest that yoga enhances individuals' ability to regulate emotional responses and adapt to environmental demands. This finding reinforces theories proposing that psychological resilience can be strengthened through intentional behavioral practices.

The observed influence of yoga frequency and duration also contributes to behavioral health theories emphasizing the cumulative effects of repeated health-promoting behaviors. The results indicate that mental health improvements increase with greater engagement in yoga activities, supporting theoretical models that view well-being as a dynamic process shaped by ongoing lifestyle choices.

Furthermore, the findings extend holistic health theories by demonstrating that interventions targeting both physical and mental processes can generate substantial psychological benefits. The study therefore provides additional evidence supporting integrated approaches to health promotion and mental health management.

### **5.6 Practical Implications:**

The practical implications of this study are significant for healthcare professionals, educational institutions, employers, policymakers, and community organizations seeking effective strategies for promoting mental health. The findings suggest that yoga represents a practical, accessible, and cost-effective intervention capable of reducing stress, anxiety, and depression among diverse populations.

Healthcare practitioners may consider incorporating yoga-based programs into preventive health initiatives and mental wellness interventions. Such programs could complement traditional treatment approaches and provide individuals with additional tools for managing psychological distress. Educational institutions may also benefit from integrating yoga sessions into student wellness programs to address academic stress, emotional challenges, and mental health concerns.

Workplace organizations can utilize yoga as part of employee wellness initiatives aimed at improving productivity, reducing occupational stress, and enhancing overall employee well-being. Regular yoga programs may contribute to healthier work environments and support long-term organizational effectiveness.

From a policy perspective, the findings support the development of community-based mental health promotion programs that include yoga and related mind-body practices. Policymakers may encourage the establishment of accessible yoga facilities, public awareness campaigns, and wellness programs that promote healthy lifestyles and psychological well-being.

Finally, individuals themselves may benefit from adopting regular yoga practice as part of a comprehensive self-care strategy. The evidence generated by this study suggests that consistent yoga participation can enhance emotional resilience, improve mental health outcomes, and contribute to a higher quality of life. Consequently, yoga may serve as an important component of preventive mental healthcare and sustainable well-being promotion.

### **6. Conclusion and Implications:**

This study examined the differences in stress, anxiety, and depression levels between yoga practitioners and non-practitioners and explored the relationship between yoga practice characteristics and mental health outcomes. The findings provide strong evidence that regular yoga participation is associated with significantly improved psychological well-being.

The comparative analysis revealed that yoga practitioners reported substantially lower levels of stress, anxiety, and depression than non-practitioners. Independent samples t-tests confirmed statistically significant differences across all three psychological indicators, supporting the study's primary hypotheses. Furthermore, correlation analysis demonstrated significant negative relationships between yoga practice frequency and psychological distress, indicating that greater engagement in yoga is associated with better mental health outcomes.

The study also found that the duration of yoga practice plays an important role in psychological well-being. Participants who had practiced yoga for longer periods reported progressively lower levels of stress, anxiety, and depression. Regression analysis further confirmed that both yoga frequency and yoga duration are significant predictors of mental health outcomes, while demographic variables such as age and gender exhibited comparatively weaker effects.

Overall, the findings suggest that yoga serves as an effective complementary approach for reducing psychological distress and promoting emotional well-being. The results reinforce the importance of regular and sustained yoga practice as a strategy for enhancing mental health and improving quality of life.

### **6.2 Academic Contributions:**

This study contributes to the growing body of literature examining the relationship between yoga and psychological well-being. Unlike many previous investigations that focused on a single mental health outcome, the present research simultaneously examined stress, anxiety, and depression within a unified comparative framework. This comprehensive approach provides a broader understanding of the influence of yoga on mental health.

The study also contributes methodologically by directly comparing yoga practitioners and non-practitioners using standardized psychological assessment tools. Such comparative evidence strengthens understanding of the differences associated with regular yoga participation and enhances the reliability of conclusions regarding its mental health benefits.

Additionally, the inclusion of yoga frequency and duration as explanatory variables extends existing knowledge regarding the intensity of engagement required to achieve psychological benefits. The findings demonstrate that mental health improvements are not merely associated with yoga participation itself but are also influenced by the consistency and longevity of practice.

From a theoretical perspective, the study supports holistic health frameworks, stress reduction theories, and mind-body intervention models by demonstrating that integrated physical, psychological, and behavioral practices contribute positively to emotional well-being. Consequently, the research enriches academic discussions regarding preventive mental healthcare and non-pharmacological interventions for psychological distress.

### **6.3 Practical Implications for Mental Health Promotion:**

The practical implications of this study highlight the value of yoga as a cost-effective, accessible, and sustainable approach to mental health promotion. Given the increasing prevalence of stress, anxiety, and depression across populations, there is a growing need for preventive interventions that can be implemented in diverse settings.

Healthcare professionals may incorporate yoga-based programs into wellness initiatives and preventive mental health services. Yoga can complement traditional treatment approaches by providing individuals with practical tools for managing stress and improving emotional regulation. Mental health practitioners may also recommend yoga as part of broader self-care and resilience-building strategies.

Educational institutions can utilize yoga programs to support students facing academic pressures, examination stress, and emotional challenges. Integrating yoga sessions into school and university wellness programs may contribute to improved concentration, emotional stability, and overall psychological well-being.

Workplace organizations may benefit from implementing yoga-based employee wellness initiatives aimed at reducing occupational stress, enhancing job satisfaction, and improving productivity. Regular workplace yoga sessions may foster healthier work environments and support employee mental health.

At the community level, yoga programs can serve as accessible interventions that encourage healthy lifestyles and psychological resilience among individuals of different age groups and backgrounds.

### **6.4 Policy Implications:**

The findings of this study have important implications for public health policy and mental health promotion strategies. Policymakers increasingly recognize the need for preventive approaches that reduce the burden of mental health disorders while minimizing healthcare costs. The evidence generated by this research suggests that yoga can contribute meaningfully to these objectives.

Government agencies and public health authorities may consider incorporating yoga and other mind-body practices into national wellness programs and mental health promotion campaigns. Community-based yoga initiatives can be developed to increase public access to preventive mental health resources, particularly in areas where professional mental health services are limited.

Educational policymakers may support the inclusion of yoga and mindfulness activities within school and university curricula to promote emotional well-being among students. Similarly, labor and occupational health authorities may encourage organizations to integrate yoga-based wellness programs into workplace health initiatives.

Public-private partnerships involving healthcare institutions, educational organizations, community centers, and yoga practitioners could facilitate the expansion of evidence-based wellness programs. Such initiatives have the potential to improve population mental health while reducing the social and economic burden associated with psychological disorders.

### **6.5 Limitations of the Study:**

Although the study provides valuable insights into the relationship between yoga practice and mental health, several limitations should be acknowledged. First, the study utilized a cross-sectional research design, which restricts the ability to establish causal relationships between yoga participation and psychological outcomes. While significant associations were identified, longitudinal research is necessary to determine causal pathways.

Second, the study relied on self-reported data, which may be subject to response bias, recall bias, and social desirability effects. Participants may have overestimated or underestimated their psychological symptoms or yoga practice behaviors.

Third, the sample was limited to a specific geographic and demographic context, which may affect the generalizability of findings to other populations. Cultural, social, and environmental factors may influence both yoga participation and mental health outcomes.

Fourth, the study did not distinguish extensively between different styles of yoga, instructor qualifications, or variations in practice intensity. These factors may influence the effectiveness of yoga interventions and warrant further examination.

Finally, additional variables such as physical activity levels, dietary habits, social support, sleep quality, and pre-existing health conditions were not included in the analysis. These factors may also contribute to psychological well-being and should be considered in future investigations.

### **6.6 Directions for Future Research:**

The limitations identified in this study provide several opportunities for future research. Longitudinal studies are recommended to examine changes in stress, anxiety, and depression over time and establish stronger causal evidence regarding the effects of yoga practice on mental health.

Future researchers may also conduct experimental and intervention-based studies that evaluate the effectiveness of specific yoga programs in reducing psychological distress. Such studies could compare different yoga styles, session durations, and practice intensities to identify the most effective approaches for mental health improvement.

Expanding the study to include larger and more diverse populations would enhance the generalizability of findings. Comparative research involving participants from different cultural, socioeconomic, and occupational backgrounds may provide additional insights into the universal and context-specific benefits of yoga.

Further investigations could examine the interaction between yoga practice and other health-related factors, including physical activity, nutrition, sleep quality, mindfulness, and social support. Understanding these relationships may contribute to the development of comprehensive wellness models.

Future research may also explore the psychological and neurobiological mechanisms underlying the observed benefits of yoga. Such investigations would deepen theoretical understanding of how yoga influences mental health and support the development of evidence-based mental health interventions.

In conclusion, this study provides substantial evidence that regular yoga practice is associated with lower levels of stress, anxiety, and depression. The findings reinforce the value of yoga as a holistic approach to mental health promotion and establish a foundation for future academic, clinical, and policy-oriented research in this important area.

**References:**

1. American Psychiatric Association. (2022). Diagnostic and statistical manual of mental disorders (5th ed., text rev.). American Psychiatric Publishing.
2. Büsing, A., Michalsen, A., Khalsa, S. B. S., Telles, S., & Sherman, K. J. (2012). Effects of yoga on mental and physical health: A short summary of reviews. *Evidence-Based Complementary and Alternative Medicine*, 2012, 1-7. <https://doi.org/10.1155/2012/165410>
3. Cramer, H., Anheyer, D., Saha, F. J., & Dobos, G. (2018). Yoga for posttraumatic stress disorder: A systematic review and meta-analysis. *BMC Psychiatry*, 18(1), 72-84. <https://doi.org/10.1186/s12888-018-1650-x>
4. Cramer, H., Lauche, R., Langhorst, J., & Dobos, G. (2013). Yoga for depression: A systematic review and meta-analysis. *Depression and Anxiety*, 30(11), 1068-1083. <https://doi.org/10.1002/da.22166>
5. Field, T. (2016). Yoga research review. *Complementary Therapies in Clinical Practice*, 24, 145-161. <https://doi.org/10.1016/j.ctcp.2016.06.005>
6. Gothe, N. P., & McAuley, E. (2015). Yoga and cognition: A meta-analysis of chronic and acute effects. *Psychosomatic Medicine*, 77(7), 784-797. <https://doi.org/10.1097/PSY.0000000000000218>
7. Gupta, N., Khera, S., Vempati, R. P., Sharma, R., & Bijlani, R. L. (2006). Effect of yoga-based lifestyle intervention on state and trait anxiety. *Indian Journal of Physiology and Pharmacology*, 50(1), 41-47.
8. Khalsa, S. B. S. (2015). Yoga for psychiatry and mental health: An ancient practice with modern relevance. *Indian Journal of Psychiatry*, 57(Suppl. 2), S239-S243. <https://doi.org/10.4103/0019-5545.161500>
9. Kirkwood, G., Rampes, H., Tuffrey, V., Richardson, J., & Pilkington, K. (2005). Yoga for anxiety: A systematic review of the research evidence. *British Journal of Sports Medicine*, 39(12), 884-891. <https://doi.org/10.1136/bjism.2005.018069>
10. Pascoe, M. C., Thompson, D. R., Jenkins, Z. M., & Ski, C. F. (2017). Mindfulness mediates the physiological markers of stress: Systematic review and meta-analysis. *Journal of Psychiatric Research*, 95, 156-178. <https://doi.org/10.1016/j.jpsy.chires.2017.08.004>
11. Riley, K. E., & Park, C. L. (2015). How does yoga reduce stress? A systematic review of mechanisms of change and guide to future inquiry. *Health Psychology Review*, 9(3), 379-396. <https://doi.org/10.1080/17437199.2014.981778>
12. Sengupta, P. (2012). Health impacts of yoga and pranayama: A state-of-the-art review. *International Journal of Preventive Medicine*, 3(7), 444-458.
13. Sharma, M. (2014). Yoga as an alternative and complementary approach for stress management. *Journal of Evidence-Based Complementary & Alternative Medicine*, 19(1), 59-67. <https://doi.org/10.1177/2156587213503344>
14. Streeter, C. C., Gerbarg, P. L., Saper, R. B., Ciraulo, D. A., & Brown, R. P. (2012). Effects of yoga on the autonomic nervous system, gamma-aminobutyric acid, and allostasis in epilepsy, depression, and post-traumatic stress disorder. *Medical Hypotheses*, 78(5), 571-579. <https://doi.org/10.1016/j.mehy.2012.01.021>
15. Telles, S., Singh, N., & Balkrishna, A. (2012). Managing mental health disorders resulting from trauma through yoga: A review. *Depression Research and Treatment*, 2012, 1-9. <https://doi.org/10.1155/2012/401513>
16. Tolahunase, M., Sagar, R., Dada, R., & Mehta, M. (2018). Impact of yoga and meditation on cellular aging in apparently healthy individuals: A prospective, open-label single-arm exploratory study. *Oxidative Medicine and Cellular Longevity*, 2018, 1-9. <https://doi.org/10.1155/2018/7928981>
17. Uebelacker, L. A., Tremont, G., Epstein-Lubow, G., Gaudiano, B. A., Gillette, T., Kalibatseva, Z., & Miller, I. W. (2010). Open trial of yoga for depression in a psychiatric outpatient population. *Complementary Therapies in Medicine*, 18(2), 68-73. <https://doi.org/10.1016/j.ctim.2010.02.003>
18. Varambally, S., & Gangadhar, B. N. (2016). Current status of yoga in mental health services. *International Review of Psychiatry*, 28(3), 233-235. <https://doi.org/10.3109/09540261.2016.1159950>
19. World Health Organization. (2022). World mental health report: Transforming mental health for all. World Health Organization. <https://www.who.int/publications/i/item/9789240049338>
20. World Health Organization. (2024). Mental health and well-being: Global status report 2024. World Health Organization.
21. Yadav, R. K., Magan, D., Mehta, M., Sharma, R., & Mahapatra, S. C. (2012). Efficacy of a short-term yoga-based lifestyle intervention in reducing stress and improving quality of life. *Indian Journal of Community Medicine*, 37(3), 165-168. <https://doi.org/10.4103/0970-0218.99912>